Astronomy & Astrophysics manuscript no. analisys © ESO 2012
December 20, 2012

arXiv:1212.3843v2 [astro-ph.EP] 18 Dec 2012

Analysis of the motion of an extrasolar planet in a binary system

E. Plávalová[1] and N.A.Solovaya[2]

[1] Astronomical Institute, Slovak Academy of Sciences, Bratislava, Slovak Republic
e-mail: `plavala@slovanet.sk`
[2] Sternberg State Astronomical Institute of the Lomonosov Moscow State University, Moscow, Russia
e-mail: `solov@sai.msu.ru`

December 20, 2012/Accepted ¡date¿

ABSTRACT

Context. More than 10% of extrasolar planets revolve in a binary or multiple stellar system. We investigated the motion of planets revolving in binary systems in the frame of the particular case of the three body problem.
Aims. We analysed of the motion an extrasolar planet revolving in a binary system by following conditions; a) a planet in a binary system revolves around one of the components (parent star), b) the distance between the star‘s components is greater than between the parent star and the orbiting planet (ratio of these two distances is a small parameter), c) the mass of the planet is smaller than the mass of the star, but is not negligible.
Methods. The Hamiltonian of the system without short periodic terms was used. Expanded in the terms of the Legendre polynomial and truncated after the second order term depending on the one angular variable. In this case the solution of the system was obtained and the qualitative analysis of motion was done. We have applied this theory to real extrasolar planets.
Results. Analysis of the possible regions of motion are presented. It is shown that the case of the stable and unstable motion of the extrasolar planets are possible. We applied our calculations to two binary systems hosting an extrasolar planet and calculated the possible values for their insufficient orbital elements. For 16 Cyg Bb, there is a region between 44° and 46° for the inclination of the planet and the value of the ascending node between 130° and 137°, or the inclination for the planet could have a value from 134° to 136° and the ascending node from 310° to 317°. For planet HD19994 b, to have a stable system, the value for the insufficient elements should be in the region of 62° to 68° for the inclination of the planet, and the value of the ascending node between 260° and 268°. Alternatively, the inclination of the planet also could have the value of 112° to 118° and the ascending node between 80° and 88°.
Conclusions. It is shown by which initial Keplerian elements the orbital motion of extrasolar planets could be stable or unstable.

Key words. Planets and satellites: general – extrasolar planets – HD 19994 Ab – 16 Cyg Bb

1. Introduction

The motion of an extrasolar planet is calculated using a frame of the particular case of the three-body problem. The planet in the binary system revolves around one of the components and the mass of the planet is much smaller than the mass of the stars, but is not negligible. The distance between the star‘s components is greater than between the parent star and the orbiting planet (the ratio of these two distances is a small parameter). The motion is considered in the Jacobian coordinate system and the invariable plane is taken as the reference plane. We used the Delaunay canonical elements L_i, G_i, and g_i (i=1 for the planet's orbit, i=2 for the star's orbit). They can be expressed through the Keplerian elements as

$$\begin{aligned} L_i &= \beta_i \sqrt{a_i}\,, \qquad G_i = L_i \sqrt{1-e_i^2}\,, \qquad H_i = G_i \, cos I_i\,, \\ l_i &= M_i\,, \qquad g_i = \omega_i\,, \qquad h_i = \Omega_i\,. \end{aligned} \tag{1}$$

Where

$$\beta_1 = k\,\frac{m_0\, m_1}{\sqrt{m_0+m_1}} = k\mu_1\,, \quad \beta_2 = k\,\frac{(m_0+m_1)\, m_2}{\sqrt{m_0+m_1+m_2}} = k\mu_2\,. \tag{2}$$

In the previous expression the notation has the usual meaning: m_0, m_2 – the masses of the stars, m_1 – the mass of the planet, k – the Gaussian constant, a_i – the semi-major axis, e_i – the eccentricity, and g_i – the argument of the perigee in the invariable plane. The eccentricities of the star's and planet's orbits can have any value from $0 < e_i < 1$. In the general case the motion is defined by the masses of components and by the six initial values of the Keplerian elements of the planet and the distant star. The formal solution of a task with the Hamiltonian without short-periodic terms was obtained in the hyperelliptic integrals by the Hamilton-Jacobi method (Orlov, 1988). Hamiltonian expanded in the terms of the Legendre polynomials and truncated after the second order terms carries the following form:

$$\begin{aligned} F &= \frac{\gamma_1}{2L_1^2} + \frac{\gamma_2}{2L_2^2} - \frac{1}{16}\gamma_3\,\frac{L_1^4}{L_2^3 G_2^3}\left[\left(1-3q^2\right)\left(5-3\eta^2\right) - \right. \\ & \left. -15\left(1-q^2\right)\left(1-\eta^2\right) cos(2g_1)\right], \end{aligned} \tag{3}$$

where the coefficients γ_1,γ_2, and γ_3 depend on mass as follows:

$$\gamma_1 = \frac{\beta_1^4}{\mu_1}\,, \qquad \gamma_2 = \frac{\beta_2^4}{\mu_2}\,, \qquad \gamma_3 = k^2\,\mu_1\,\mu_2\,\frac{\beta_2^6}{\beta_1^4}\,, \tag{4}$$

the parameter:

$$\eta = \sqrt{1-e_1^2}\,, \tag{5}$$

and the cosine of the angle between the plane of the planet orbit and the plane of the distant star's orbit

$$q = \frac{c^2 - G_1^2 - G_2^2}{2\,G_1\,G_2} = cos\,(i_1 + i_2)\;. \tag{6}$$

c is the constant of the angular momentum, and g_1 is the argument of the perigee of the planet orbit in the invariable plane.
The Hamilton's equation truncated after the second order terms depends on the one angular variable g_1 only. It is necessary to investigate these equations

$$\frac{dG_1}{dt} = \frac{\partial F}{\partial g_1}\,, \qquad \frac{dg_1}{dt} = -\frac{\partial F}{\partial G_1}\,. \tag{7}$$

In this approximation the task is integrated. This solution we have used for intermediate motion. So the semi-major axis a_1 and a_2 have no the secular terms; they are restricted. But the eccentricity of the planet's orbit can be changed significantly, almost as much as 1. Then the system will be come unstable. The change of the eccentricity of an extrasolar planet is described by the expression $e_1 = \sqrt{1-\xi}$. Generally, the dependence between ξ and t is defined by the equation (Orlov,1988):

$$\frac{1}{12}\overline{G}_2^2 \int_{\xi_1}^{\xi} \frac{\mathrm{d}\xi}{\sqrt{\Delta}} = \frac{B_3}{A_1} + \frac{1}{16}\frac{\gamma m''^2}{\left(1-e_2^2\right)^{\frac{3}{2}}} n_1\,(t-t_0)\,. \tag{8}$$

Where $m" = n_2/n_1$ and n_1, n_2 are the mean motions of the planet and star, $\xi = G_2^2/L_1^2$, $\gamma = m_2/m_0 + m_1 + m_2$ while, A_1 and B_3 are the constants of integration.
On the left side of the equation (8) under the integral in the denominator is the square root from the polynomial of the fifth order. This polynomial can be presented as the product of the two polynomials of the second and the third orders, $\Delta = f_2(\xi)f_3(\xi)$, where $\xi = 1 - e_1^2$, and e_1 is the eccentricity of the extrasolar planet's orbit. The establishment of the regions of the motion are possible when the roots of the equations $f_2(\xi) = 0$, $f_3(\xi) = 0$ are found and the signs of the functions defined.

2. Equations of the second and third order

We investigated the roots of the equations $f_2(\xi) = 0$ and $f_3(\xi) = 0$ with following forms:

$$f_2(\xi) = \xi^2 - 2\left(\overline{c}^2 + 3\overline{G}_2^2\right)\xi + \left(\overline{c}^2 - \overline{G}_2^2\right)^2 + \frac{2}{3}\left(10 + A_3\right)\overline{G}_2^2 \tag{9}$$

and

$$\begin{aligned} f_3(\xi) = {} & \xi^3 - \left(2\overline{c}^2 + \overline{G}_2^2 + \frac{5}{4}\right)\xi^2 + \left[\frac{5}{2}\left(\overline{c}^2 + \overline{G}_2^2\right) + \right. \\ & \left. + \left(\overline{c}^2 - \overline{G}_2^2\right)^2 - \frac{1}{6}\overline{G}_2^2\left(10 + A_3\right)\right]\xi - \frac{5}{4}\left(\overline{c}^2 - \overline{G}_2^2\right)^2\,. \end{aligned} \tag{10}$$

Where

$$\overline{c} = \frac{c}{L_1}\,, \qquad \overline{G}_2 = \frac{G_2}{L_1} = \frac{\beta_2}{\beta_1}\sqrt{\frac{a_2\left(1-e_2^2\right)}{a_1}}\,, \quad \overline{G}_2 > 1\,, \tag{11}$$

and

$$A_3 = 2 - 6\eta_0^2 q_0^2 - 6\left(1-\eta_0^2\right)\left[2 - 5\left(1-q_0^2\right)sin^2 g_{10}\right]\,. \tag{12}$$

We noted

$$\eta_0 = \sqrt{1-e_{10}^2}\,, \tag{13}$$

where, e_{10} is the initial value of the eccentricity of the planet, q_0 is the initial value of the cosine of the mutual inclination between the orbits of the planet and it's distant star and g_{10} is the initial value of the argument of the perigee of the planet's orbit in the invariable plane.
So the orbit is assumed to be elliptic, $0 < e_1 < 1$, then $0 < \xi < 1$. Value of $\xi = 1$ corresponds on the circular motion.
For establishment of the boundaries of the regions of the possible motion it is necessary to find the roots the equations (9) and (10) and to define the signs of the function in the interval between the roots.

2.1. The equation of the second order

We rewrote equation of the second order (9) in the following form:

$$f_2(\xi) = \xi^2 - 2\left(\overline{c}^2 + 3\overline{G}_2^2\right)\xi + \left[-1 + 2\left(\overline{c}^2 + 3\overline{G}_2^2\right) + \frac{2}{3}\overline{G}_2^2\overline{h}\right].\tag{14}$$

Then defined $\overline{h}$ as difference between values A_3 and A_{3crit}, where

$$\overline{h} = A_3 - A_{3crit}\,, \tag{15}$$

and

$$\begin{aligned} 2\overline{G}_2^2\overline{h} = {} & 3(1-\eta^2)\left[1 - 8\overline{G}_2^2 - 4\overline{G}_2 q\eta - \eta^2 + \right. \\ & \left. + 20\overline{G}_2^2\left(1-q^2\right)sin^2 g_1\right]\,. \end{aligned} \tag{16}$$

We named A_{3crit} the meaning of the constant A_3 for which the square equation has the root equal to one. This meaning is found in the left part of the $f_2(\xi) = 0$, $\xi = 1$. We note the roots of the equation (14) as ϵ_1 and ϵ_2, and $\epsilon_1 < \epsilon_2$. The coefficient by ξ is always less than zero. This equation has no negative roots and the free term is also greater then 0. The roots are

$$\epsilon_{1,2} = \left(\overline{c}^2 + 3\overline{G}_2^2\right) \mp \sqrt{\left(\overline{c}^2 + 3\overline{G}_2^2 + 1\right)^2 - \frac{2}{3}\overline{G}_2^2\overline{h}}\,, \tag{17}$$

and $\epsilon_2 >> 1$ always. Derivation $\frac{d\epsilon_1}{d\overline{h}}$ is always positive. So ϵ_1 is a growing function of $\overline{h}$. When $\overline{h} = 0$ then $\epsilon_1 = 1$, we conclude for $\overline{h} < 0$ then $\epsilon_1 < 1$, and for $\overline{h} > 0$ then $\epsilon_1 > 1$. For the definition of the branch of the parabola we found the extreme of the function in the point which is valid $\overline{\xi} = \overline{c}^2 + 3\overline{G}_2^2$. The value $f_2(\overline{\xi}) < 0$ and the branches of this parabola are directed up.

2.2. Equation of the third order

We rewrote equation (10) using parameter $\overline{h}$ in the form:

$$\begin{aligned} f_3(\xi) = {} & \xi^3 - \frac{1}{4}\left(5 + 8\overline{c}^2 + 4\overline{G}_2^2\right)\xi^2 + \left[\frac{1}{4} + 2\overline{c}^2 + \overline{G}_2^2 + \right. \\ & \left. + \frac{5}{4}\left(\overline{c}^2 - \overline{G}_2^2\right)^2 - \frac{1}{6}\overline{G}_2^2\overline{h}\right]\xi - \frac{5}{4}\left(\overline{c}^2 - \overline{G}_2^2\right)^2\,. \end{aligned} \tag{18}$$

Then multiplied this equation by 4 and identified in its left part the item $\xi f_2(\xi)$. Then the equation (18) can be rewrote in the following form:

$$5\,(1-\xi)\left[\left(\overline{c} + \overline{G}_2\right)^2 - \xi\right]\left[\xi - \left(\overline{c} - \overline{G}_2\right)^2\right] - \xi f_2(\xi) = 0\,. \tag{19}$$

For the establishment of the qualitative characteristics of the motion, it is necessary to define how the roots of the equation of the third order are located in relation to the points of the axes:

$$\xi = 0, \qquad \xi = \epsilon_1, \qquad \xi = 1, \qquad and \qquad \xi = \epsilon_2. \tag{20}$$

1. If $\overline{h} < 0$ then we have:
   (a) $\xi = 0, \qquad f_3(0) < 0.$
   (b) $\xi = \epsilon_1, \qquad f_3(\epsilon_1) > 0.$
   (c) $\xi = 1, \qquad f_3(1) > 0.$
   (d) $\xi = \epsilon_2, \qquad f_3(\epsilon_2) > 0.$
   From Sturm theorem (e.g. Dörrie 1965) it follows that, between 0 and ϵ_1 lay at least one root of the third order equation.
2. If $\overline{h} > 0$ then we have:
   (a) $\xi = 0\,, \qquad f_3(0) \leqslant 0.$
   (b) $\xi = 1\,, \qquad f_3(1) < 0.$
   (c) $\xi = \epsilon_1\,, \qquad f_3(\epsilon_1) < 0.$
   (d) $\xi = \epsilon_2\,, \qquad f_3(\epsilon_2) > 0.$

The behaviours of the roots of the second and third order equations depend on the sign of $\overline{h}$ and valid:

1. for $\overline{h} < 0$ is valid $0 \leqslant \epsilon_3 < \epsilon_1 < 1 < \epsilon_4 < \epsilon_5 < \epsilon_2$.
2. for $\overline{h} > 0$ is valid $0 \leqslant \epsilon_3 < \epsilon_4 < 1 < \epsilon_1 < \epsilon_5 < \epsilon_2$.

We have two roots which are less then 1, therefore we identified the roots in ascending order as $\xi_1, \xi_2, \xi_3, \xi_4, \xi_5$. This means that e_1 can change in limits from $e_{1min} = 1 - \xi_2$ to $e_{1max} = 1 - \xi_1$. We showed the behaviour of these functions for the hypothetical value $\overline{c}$, $\overline{G}_2$ and $\overline{h}$ on the Figure (2.2) for $\overline{h} < 0$ and on Figure (2.2) for $\overline{h} > 0$.



Fig. 1. Behaviour of the functions $f_2(\xi)$ (solid line) and $f_3(\xi)$ (dashed line) for $\overline{h} > 0$. The left-side graph shows an overview while the right-side graph shows the detail between zero and two.



Fig. 2. Behaviour of the functions $f_2(\xi)$ (solid line) and $f_3(\xi)$ (dashed line) for $\overline{h} < 0$. The left-side graph shows an overview while the right-side graph shows the detail between zero and two.

3. The investigation of the value $\overline{\text{h}}$

Consider the value $\overline{h}$, which has the following form:

$$\overline{h} = \frac{3(1-\eta^2)}{2\overline{G}_2^2}\Big[1 - 8\overline{G}_2^2 - 4\overline{G}_2 q\eta - \eta^2 + \\ +20\overline{G}_2^2\left(1-q^2\right)sin^2 g_1\Big]\,. \qquad (21)$$

When the value $g_1 = 0$ then value $\overline{h}$ reaches minimum

$$\overline{h}_{min} = \frac{3(1-\eta^2)}{2\overline{G}_2^2}\left[1 - 8\overline{G}_2^2 - 4\overline{G}_2 q\eta - \eta^2\right]. \qquad (22)$$

The value of the $\overline{h}_{min}$ is always negative. In this case we have the two roots valued less then 1, the one root of the equation of the third order and one root of the square equation. If $g_1 = \pi/2$ then $\overline{h}$ has a maximum value

$$\overline{h}_{max} = \frac{3(1-\eta^2)}{2\overline{G}_2^2}\left[1 - \eta^2 - 4\overline{G}_2 q\eta + 12\overline{G}_2^2 - 20\overline{G}_2^2 q^2\right]. \qquad (23)$$

The value of $\overline{h}_{max}$ could have both a positive and negative value. For the case $g_1 = \pi/2$ we rewrote equations (9) and (10) in the following form:

$$f_2(\xi) = \left(\xi - \eta_0^2\right)\left(\xi - \eta_0^2 - 4\overline{G}_2\eta_0 q_0 + 8\overline{G}_2^2\right) + \qquad (24)$$

$$+4\overline{G}_2^2\left(1-5q_0^2\right)\left(1-\eta_0^2\right) + 16\overline{G}_2^2\,, \qquad (25)$$

and

$$f_3(\xi) = \left(\xi - \eta_0^2\right)\Bigg[\left(\xi - \frac{5}{4}\right)\left(\xi - \eta_0^2\right) - \\ -3\overline{G}_2^2\xi + \overline{G}_2\eta_0 q_0\left(5 - 4\xi\right) + 5\overline{G}_2^2 q_0^2\Bigg]\,. \qquad (26)$$

When one of the roots of the equation (26) carries the value of $\xi = \eta_0^2$, the other roots have to satisfy the equation

$$\left(\xi - \frac{5}{4}\right)\left(\xi - \eta_0^2\right) - 3\,\overline{G}_2^2\,\xi + \overline{G}_2\,\eta_0\,q_0\left(5 - 4\xi\right) + 5\,\overline{G}_2^2\,q_0^2 = 0\,. (27)$$

For $\overline{h}_{max}$ the starting value $\eta^2 = \eta_0^2$ is one of the boundary limits for change in the value of ξ. We should establish that $\xi = \eta_0^2$ is the least root and the second root has a value less than 1. We therefore substitute the value $\xi = \eta_0^2$ in (27). If the obtained expression is negative, then η_0^2 should become the second root in equation (26), if positive – then it should become the least root. The left part of the equation (27) for $\xi = \eta_0^2$ is

$$2\left[5q_0^2 - 3\eta_0^2 + \frac{1}{\overline{G}_2}\eta_0 q_0\left(5 - 4\eta_0^2\right)\right]. \qquad (28)$$

Equating this expression to zero and solving the obtained equation concerning q_0, we find the value of q_0 of which η_0^2 is the root of expression(28). So

$$q = \frac{\eta_0\left[4\eta_0^2 - 5 \mp \sqrt{60\overline{G}_2^2 + \left(5 - 4\eta_0^2\right)^2}\right]}{10\overline{G}_2}\,. \qquad (29)$$

We denote the solution of equation (29) as q_{01} for the minus sign before the root term and the q_{02} for plus sign. If the value of q_0 lays inside the region of $q_{01} < q_0 < q_{02}$, expression (28) is negative, and the value of the root lays between the other two roots. If either of the conditions $q_0 < q_{01}$ or $q_0 > q_{02}$ is valid, then $\xi = \eta_0^2$ is the least root of equation (27). In this case $e_{1_{\max}} = \sqrt{1 - \eta_0^2}$ and the maximum value of the eccentricity of the planet's orbit can not exceed the initial value of eccentricity. The orbit of the extrasolar planet may be stable.

Table 1. Initial orbital elements of the system 16 Cyg.

	star 16 Cyg A	planet 16 Cyg Bb
Mass [M_{Sun}]	1.53	
Mass [M_{Jup}*sin i_1]		1.68±0.07
Semi-major axis [AU]	754.53	1.68±0.07
Eccentricity	0.863	0.689±0.011
Inclination	135.44°	
Ascending node	313.44°	
Perigee	26.6°	83.4±2.1°
Period	13512.7 years	799.5 days

Notes. Mass of the parent star 16 Cyg B is 1.01 ± 0.04M_{Sun}. We used the planet's semi-major axis value 1.6923 AU for the calculations.

Table 2. Our proposal for a full set of orbital elements for planet 16 Cyg Bb.

	First possibility	Second possibility
Mass[M_{Jup}]	2.38±0.04	2.38±0.04
Semi-major axis [AU]	1.693	1.693
Eccentricity	0.689±0.011	0.689±0.011
Inclination	45°±1°	135°±1°
Ascending node	133.5°±3.5°	313.5°±3.5°
Perigee	83.4±2.1°	83.4±2.1°
Period[day]	799.5	799.5

4. Application of the theory on real extrasolar planets

The knowledge of the six pairs of Keplerian elements of the orbits of the system allow us to investigate the character of the evolution of the planet's orbit and the possible conditions of stability. They may be presented by the orbital parameters, which we obtain from the analytical theory. They are – the angle of the mutual inclination between orbits of the planet and star, the angular moment of the star and the maximum value of the eccentricity of the planet's orbit. The growth of the eccentricity of the orbit could lead to the destruction of the orbit in perigee from tidal forces. In the catalogue of extrasolar planets there is a lack of data regarding the insufficient elements for the longitude of the node and the inclination. Our theory allows for a range of possible values for the insufficient elements. In our application of the theory, we have selected two binary stellar systems, with a planet revolving around one of the components. The first is system 16 Cyg with an interesting planet 16 Cyg Bb, which has a perigee close to 90°. The second system is HD19994.

4.1. 16 Cyg

16 Cyg A (HD 186408) and 16 Cyg B (HD 186427) are both members of a well-known wide binary system with stars spectral types G1.5V and G3V. For the calculations, we used orbital elements used by Hauser and Marcy (1998), which were published in the Sixth Catalogue of Orbits of Visual Binary Stars (Hartkopf and Mason 2001). The semi-major axis was calculated to be 754.53 AU using the values 21.41 pc (Fuhrmann et. al 1998) for distance and 35.242″ for angular separation. We made a revision calculation for the semi-major axis using the third Keplerian law and the value of a 13512.7 year period (Hartkopf and Mason 2001). We reached a value of 774.06 AU. There is an obvious difficulty in determining orbital parameters for periods greater than 10,000 years. Only a small fraction of the orbit has transpired since astrometric measurements were first obtained in the early 1800s. So, we used the value 754.53 AU, derived from the distance, for the following calculations because we considered the determination period less precise than the determination of the values for angular separation and distance.

The secondary star in the 16 Cyg binary system, 16 Cyg B (parent star), with a mass of 1.01±0.04 M_{Sun} (Fuhrmann et. al 1998), is known to host a giant gas planet 16 Cyg Bb with taxonomy class 2J0.2W7 (Plávalová 2012). The extrasolar planet has a minimum mass $M_{Jup} * sin(i) = 1.68 \pm 0.07 M_{Jup}$ in an orbit with high eccentricity $e = 0.689$ (Wittenmyer et. al 2007). The starting conditions for investigating this planet, we took from The Extrasolar Planets Encyclopaedia (Schneider et. al 2012). The values of the inclination and ascending note for the planet are unknown. In Table (1). are shown the orbital elements for the planet and distant star.

We made a revision calculation for the semi-major axis of the planet, using the third Keplerian law and values for the orbital period of the planet, which is 799.5 days and published masses for the planet and mother star (Table 1). We calculated it to be $a_1 = 1.6923$ AU. The mass of the planet is defined as minimum mass and if we varied this mass to the value of 5 M_{Jup} we would get a value of 1.6941 AU. According to the results of previous calculations, we decided to use the value 1.6923 AU for the semi-major axis and 1.68 M_{Jup} for the mass of the planet for our calculations.

We varied the elements of the unknown values for 16 Cyg Bb in our calculations; the value for the ascending node from 0° to 360°, 1° at a time, and the value for the inclination from 0° to 180°, also for each degree. We recognized the pairs of the elements which when applied to the planet, its motion would be unstable, and the others possibly stable. For example for the values,

$$i_1 = 44°, \text{ and } \Omega_1 = 134°,$$

the value $\bar{h}$ is near to $\bar{h}_{\max}$ ($\bar{h} = -0.5696$, $\bar{h}_{max} = -0.5698$). We can use the conditions of stability $q_0 < q_{01}$ ($q_0 = -0.9999$, $q_{01} = -0.5614$), and the expression (28)> 0 and the maximum value of the eccentricity is 0.689. With these elements the planet's orbit would be stable. Contrariwise, for example, for values,

$$i_1 = 40°, \text{ and } \Omega_1 = 20°,$$

there is a difference between values $\bar{h}$ and $\bar{h}_{\max}$ ($\bar{h} = 0.2129$, $\bar{h}_{max} = -0.5721$), $q_0 > q_{01}$ ($q_0 = -0.3670$, $q_{01} = -0.5603$), and $e_{1_{\max}} = 0.9295$. With these elements the planet's orbit would be unstable.

We found the pairs of the unknown elements, where the one root of the second and third order equations have identical values. The following are the results we obtained (Table 2). We proposed for a stable orbit, these two possible sets of values for the unknown elements; the inclination for the planet could have a value between 44° and 46° and the ascending node from 130° to 137°, or the inclination of the planet could have a value between 134° and 136° and the ascending node from 310° to 317°. These values for a stable orbit correspond to the mass of a planet equal to 2.38±0.04 M_{Jup} (for $i_1 = 44°$ is 2.42 M_{Jup} and for $i_1 = 46°$ is 2.34 M_{Jup}) calculated from a catalogue value of $M_{Jup} * sin(i) = 1.68$. We recalculated the planet's semi-major axis with this new mass applied through Keplerian law. The new value is 1.693 AU. In Table (2) is shown a proposal for two full sets of orbital elements for this planet.

Table 3. Initial orbital elements of the system HD19994.

	star HD19994	planet HD19994 b
Mass [M_{Sun}]	0.37	
Mass [M_{Jup}*sin i_1]		1.68
Semi-major axis[AU]	151.51	1.42
Eccentricity	0.26	0.3±0.04
Inclination	114.1°	
Ascending node	84.13°	
Perigee	247.74°	41±8°
Period	1420 years	535.7±3.1 days

Notes. Mass of the parent star HD19994 is 1.34M_{Sun}. We used the planet's semi-major axis value 1.4273 AU for the calculations.

4.2. HD 19994

The binary stellar system HD 19994 (94 Ceti, ADS 2406 AB) contains an A component; a yellow-white dwarf with a mass of 1.34 M_{Sun} and a B component; a red dwarf with a mass of 0.37 M_{Sun}. For the distant star we used Keplerian elements published by Hale (1994). To define the semi-major axis we used two methods. Firstly, we derived this value using stellar parallax, where the distance was 22.38 pc (Schneider 2012) and 6.77″ for the angular separation (Hale 1994) and resulted in a semi-major axis of 151.51 AU. Secondly, with the application of Keplerian law, using the values from Table (3), we calculated the value to be 151.37 AU. The difference between these two values is negligible we applied the value 151.51 AU in the following calculations.

The planet HD 19994 b was discovered in 2000 (Queloz 2001) and is orbiting an A component. The taxonomy class is 2J0.2G3 (Plávalová 2012) and the minimum mass is 1.68 M_{Jup}. This planet rotates with a semi-major axis of 1.42 AU with quite a high eccentricity 0.3 ± 0.04. For our calculations we used the orbital elements published by Mayor et. al. (2004). In Table (3) the initial Kepler orbital elements for the planet and its distant star are shown. As with the first system, we revised the calculation for the semi-major axis of the planet using Keplerian law. With the values listed in Table (3), we valued the planet's semi-major axis at a_1=1.4273 AU. If we varied the mass of the planet to 5 M_{Jup} we would get a value of 1.4284 AU. We decided to use the value 1.4273 AU for the semi-major axis and 1.68 M_{Jup} for the mass of the planet in our calculations.

We varied the values for the inclination and ascending node for the planet in our calculations, as in the first system. We obtained the following results. The stable orbit can have the following value of elements; the planet's inclination could have a value from 62° to 68° and ascending node between 260° and 268°. Alternatively, the inclination for the planet could have a value from 112° to 118° and the ascending node between 80° and 88° (Table 4). These values for a stable orbit correspond to the mass of a planet equal to 1.855±0.045 M_{Jup} (for $i_1 = 62°$ is 1.90 M_{Jup} and for $i_1 = 68°$ is 1.81 M_{Jup}) calculated from a catalogue value of $M_{Jup} * sin(i) = 1.68$. We recalculated the planet's semi-major axis with this new mass applied through Keplerian law. The new value is 1.427 AU. In Table (4) is shown a proposal for two full sets of orbital elements for this planet.

5. Conclusion

We have shown that an extrasolar planet revolving in a binary system around one of the components, may rotate on a stable or unstable orbits. The conditions for a stable motion depend on the orbital parameters, which can be calculated from the formulas of the previous sections. There are the angle of the mutual inclination between orbits, the angular momentum of the distant star and the maximum value of the eccentricity of the planet's orbit. When the value of the planet's eccentricity is close to one, in a perigee, the tidal forces of the star could destroy the planet. We have suggested the possible values for the insufficient elements with which the orbit of the planet would remain stable. For 16 Cyg Bb there are a two regions of stability. Firstly, from 44° to 46° for the inclination of the planet and for the ascending node, values from 130° to 137°. Secondly, the planetal inclination value could be between 134° and 136° and ascending node from 310° to 317°. We proposed the planet's mass to be 2.38±0.04 M_{Jup} and the value of its semi-major axis equal to 1.693 AU. For the second system HD19994, the values which would suggest a stable motion are in the region of 62° to 68° for planetal inclination and 260° to 268° for its ascending node. Alternatively, we could suggest 112° to 118° for the inclination and 80° to 88° for the ascending node. For the planet's mass we proposed the value of 1.855±0.045 M_{Jup} and for the planet's semi-major axis a value of 1.427 AU.

Table 4. Our proposal for a full set of orbital elements for planet HD19994 b.

	First possibility	Second possibility
Mass[M_{Jup}]	1.855±0.045	1.855±0.045
Semi-major axis [AU]	1.427	1.427
Eccentricity	0.3±0.04	0.3±0.04
Inclination	65°±3°	115°±3°
Ascending node	264°±4°	84°±4°
Perigee	41°±8	41°±8
Period[day]	535.7±3.1	535.7±3.1

Acknowledgements. This work was supported by the Slovak Research and Development Agency under the contract No. APVV-0158-11.

References

Fuhrmann, K., Pfieffer, M.J. and Bernkopf, J. 1998, A&A, 942

Hauser, H.M and Marcy G.W 1998, PASP 111, 321

Dörrie, H. "Sturm's Problem of the Number of Roots." §24 in 100 Great Problems of Elementary Mathematics: Their History and Solutions. New York: Dover, pp. 112-116

Hale, A. 1994,Astron.J., 107, 1, 306

Hartkopf and Mason 2001,Six Catalog of Orbits of Visual Binary Stars.

Mayor, M., Udry, S., Naef, D., Queloz, D., Santos, N.C., and Burnet, M. 2004, A&A 415, 391

Orlov, A.A., Solovaya, N.A. 1988, in The Few Body Problem, ed: M.J.Valtonen, Kluwer Acad. Publish., Dordrecht, 243

Plávalová, E. 2012, Astrobiology, 12, 4, 361

Queloz, D., Mayor, M. 2001, The Messenger, 105, 1-7

Schneider, J., Dedieu, C., Le Sidaner, P., Savalle, R., and Zolotukhin,I. 2012, A&A, 532:A79

Solovaya, N.A. and Pittich, E.M. 2004, Contrib. Astron. Obs. Skalnaté Pleso 34, 105

Wittenmyer, R.A. and Cochran, W.D. 2007, ApJ, 654,625